

PERTURBATION ANALYSIS FOR THE GENERALIZED INVERSES WITH PRESCRIBED IDEMPOTENTS IN BANACH ALGEBRAS

JIANBING CAO^{1,2} and YIFENG XUE¹

¹Department of Mathematics
East China Normal University
Shanghai 200241
P. R. China
e-mail: yfxue@math.ecnu.edu.cn

²Department of Mathematics
Henan Institute of Science and Technology
Xinxiang, Henan, 453003
P. R. China
e-mail: caocjb@163.com

Abstract

In this paper, we first study the perturbations and expressions for the generalized inverses $a_{p,q}^{(2)}$, $a_{p,q}^{(1,2)}$, $a_{p,q}^{(2,l)}$, and $a_{p,q}^{(l)}$ with prescribed idempotents p and q . Then, we investigate the general perturbation analysis and error estimate for some of these generalized inverses when p , q and a also have some small perturbations.

2010 Mathematics Subject Classification: 15A09, 46L05.

Keywords and phrases: gap function, idempotent element, generalized inverse, perturbation.

Communicated by Jong-Kwang Yoo.

Received December 3, 2012

1. Introduction

Let \mathcal{R} be a unital ring and let \mathcal{R}^\bullet denote the set of all idempotent elements in \mathcal{R} . Given $p, q \in \mathcal{R}^\bullet$. Recall that an element $a \in \mathcal{R}$ has the (p, q) -outer generalized inverse $b = a_{p,q}^{(2)} \in \mathcal{R}$ if $bab = b$, $ba = p$, and $1 - ab = q$. If $b = a_{p,q}^{(2)}$ also satisfies the equation $aba = a$, then we say a has the (p, q) -generalized inverse b , in this case, written $b = a_{p,q}^{(1,2)}$. If an outer generalized inverse with prescribed idempotents exists, it is necessarily unique (cf. [6]). According to this definition, obviously, we see that the Moore-Penrose inverses in a C^* -algebra and (generalized) Drazin inverses in a Banach algebra can be expressed by some (p, q) -outer generalized inverses (cf. [1, 5, 6]).

Based on some results of Djordjević and Wei in [6], Ilic et al. gave some equivalent conditions for the existence of the (p, q) -outer generalized inverse in a Banach algebra in [5]. But in our recent paper [1], we find that Theorem 1.4 of [5] is wrong. In [1], we first present a counter-example to [5, Theorem 1.4], then based on our counter-example, we define a new type of generalized inverse with prescribed idempotents in a Banach algebra as follows:

Definition 1.1 (see [1]). Let $a \in \mathcal{A}$ and $p, q \in \mathcal{A}^\bullet$. An element $b \in \mathcal{A}$ satisfying

$$bab = b, \quad R_r(b) = R_r(p), \quad K_r(b) = R_r(q),$$

will be called the (p, q, l) -outer generalized inverse of a , written as $a_{p,q}^{(2,l)} = b$.

In addition, if $a_{p,q}^{(2,l)}$ satisfies $a = aa_{p,q}^{(2,l)}a$, we call $a_{p,q}^{(2,l)}$ is the (p, q, l) -generalized inverse of a , denoted by $a_{p,q}^{(l)}$.

Perturbation analysis of the generalized inverses is very important in both theory and applications. In recent years, there are many fruitful results concerning the perturbation analysis for various types generalized inverses of operators on Hilbert spaces or Banach spaces. The concept of stable perturbation of an operator on Hilbert spaces and Banach spaces is introduced by Chen and Xue in [1]. Later, the notation is generalized to the set of Banach algebras by the second author in [12] and to the set of Hilbert C^* -modules by Xu et al. in [15]. Using the notation “stable perturbation”, many important results in perturbation analyses for Moore-Penrose inverses on Hilbert spaces and Drazin inverses on Banach spaces or in Banach algebras have been obtained. Please see [2, 3, 4, 12, 13, 14] for detail.

Let X, Y be Banach spaces over complex field \mathbb{C} . Let T (resp., S) be a given closed space in X (resp., Y). Let A be a bounded linear operator from X to Y such that $A_{T,S}^{(2)}$ exists. The perturbation analysis of $A_{T,S}^{(2)}$ for small perturbation of T, S , and A has been done in [7]. Motivated by some recent results, concerning the perturbation analysis for the generalized inverses of operators, in this paper, we mainly study the perturbations and expressions for various types of generalized inverses with prescribed idempotents in Banach algebras. We first consider the stable perturbation characterizations for $a_{p,q}^{(2)}$, $a_{p,q}^{(1,2)}$, $a_{p,q}^{(2,l)}$, and $a_{p,q}^{(l)}$ with prescribed idempotents p and q . Then, by using stable perturbation characterizations, we can investigate the general perturbation analysis and error estimate for some of these generalized inverses when p, q , and a also have some small perturbations. The results obtained in this paper extend and improve many recent results in this area.

2. Preliminaries

In this section, we give some notations in this paper, we also list some preliminary results, which will be frequently used in our main

sections. Throughout the paper, \mathcal{A} is always a complex Banach algebra with the unit 1.

Let $a \in \mathcal{A}$. If there is $b \in \mathcal{A}$ such that $aba = a$ and $bab = b$, then a is called to be generalized invertible and b is called the generalized inverse of a , denoted by $b = a^+$. Let $Gi(\mathcal{A})$ denote the set of all generalized invertible elements in $\mathcal{A} \setminus \{0\}$. Let \mathcal{A}^\bullet denote the set of all idempotent elements in \mathcal{A} . If $a \in Gi(\mathcal{A})$, then a^+a and $1 - aa^+$ are all idempotent elements. For $a \in \mathcal{A}$, set

$$\begin{aligned} K_r(a) &= \{x \in \mathcal{A} \mid ax = 0\}, & R_r(a) &= \{ax \mid x \in \mathcal{A}\}; \\ K_l(a) &= \{x \in \mathcal{A} \mid xa = 0\}, & R_l(a) &= \{xa \mid x \in \mathcal{A}\}. \end{aligned}$$

Clearly, if $p \in \mathcal{A}^\bullet$, then \mathcal{A} has the direct sum decompositions

$$\mathcal{A} = K_r(p) \dot{+} R_r(p) \quad \text{or} \quad \mathcal{A} = K_l(p) \dot{+} R_l(p).$$

The following useful and well-known lemma can be easily proved.

Lemma 2.1. *Let $x \in \mathcal{A}$ and $p \in \mathcal{A}^\bullet$. Then*

- (1) $K_r(p)$ and $R_r(p)$ are all closed and $K_r(p) = R_r(1 - p)$, $R_r(p) \mathcal{A} \subset R_r(p)$;
- (2) $px = x$ if and only if $R_r(x) \subset R_r(p)$ or $K_l(p) \subset K_l(x)$;
- (3) $xp = x$ if and only if $K_r(p) \subset K_r(x)$ or $R_l(x) \subset R_l(p)$.

We list some of the necessary and sufficient conditions for the existence of $a_{p,q}^{(2,l)}$ in the following lemma, which will be frequently used in the paper. Here, we should indicate that $a_{p,q}^{(2,l)}$ is unique if it exists. Please, see [1] for the proofs and more information.

Lemma 2.2. *Let $a \in \mathcal{A}$ and $p, q \in \mathcal{A}^\bullet$. Then the following statements are equivalent:*

- (1) $a_{p,q}^{(2,l)}$ exists;
- (2) there exists $b \in \mathcal{A}$ such that $bab = b$, $R_r(b) = R_r(p)$, and $K_r(b) = R_r(q)$;
- (3) $K_r(a) \cap R_r(p) = \{0\}$ and $\mathcal{A} = aR_r(p) \dot{+} R_r(q)$;
- (4) there exists $b \in \mathcal{A}$ satisfying $b = pb$, $p = bap$, $b(1-q) = b$, $1-q = (1-q)ab$;
- (5) $p \in R_l((1-q)ap) = \{x(1-q)ap \mid x \in \mathcal{A}\}$ and $1-q \in R_r((1-q)ap)$;
- (6) there exist some $s, t \in \mathcal{A}$ such that $p = t(1-q)ap$, $1-q = (1-q)aps$.

The following lemma gives some equivalent conditions about the existence of $a_{p,q}^{(l)}$. See [1] for more information.

Lemma 2.3. *Let $a \in \mathcal{A}$ and $p, q \in \mathcal{A}^\bullet$. Then the following conditions are equivalent:*

- (1) $a_{p,q}^{(l)}$ exists, i.e., there exists some $b \in \mathcal{A}$ such that
- $$aba = a, \quad bab = b, \quad R_r(b) = R_r(p), \quad K_r(b) = R_r(q),$$
- (2) $\mathcal{A} = R_r(a) \dot{+} R_r(q) = K_r(a) \dot{+} R_r(p)$,
- (3) $\mathcal{A} = aR_r(p) \dot{+} R_r(q)$, $R_r(a) \cap R_r(q) = \{0\}$, $K_r(a) \cap R_r(p) = \{0\}$.

Let X be a complex Banach space. Let M, N be two closed subspaces in X . Set

$$\delta(M, N) = \begin{cases} \sup \{\text{dist}(x, N) \mid x \in M, \|x\| = 1\}, & \text{if } M \neq \{0\}, \\ 0 & \text{if } M = \{0\}, \end{cases}$$

where $\text{dist}(x, N) = \inf \{\|x - y\| \mid y \in N\}$. The gap $\hat{\delta}(M, N)$ of M, N is given by $\hat{\delta}(M, N) = \max \{\delta(M, N), \delta(N, M)\}$. For convenience, we list

some properties about $\delta(M, N)$ and $\hat{\delta}(M, N)$, which come from [9] as follows:

Proposition 2.4 ([9]). *Let M, N be closed subspaces in a Banach space X . Then*

- (1) $\delta(M, N) = 0$ if and only if $M \subset N$;
- (2) $\hat{\delta}(M, N) = 0$ if and only if $M = N$;
- (3) $\hat{\delta}(M, N) = \hat{\delta}(N, M)$;
- (4) $0 \leq \delta(M, N) \leq 1, 0 \leq \hat{\delta}(M, N) \leq 1$.

3. Stable Perturbations for the (p, q) -Generalized Inverses

Let $a \in Gi(\mathcal{A})$ and let $\bar{a} = a + \delta a \in \mathcal{A}$. Recall from [13] that \bar{a} is a stable perturbation of a if $R_r(\bar{a}) \cap K_r(a^+) = \{0\}$. Obviously, we can define the stable perturbation for various kind of generalized inverses. In this section, we concern the stable perturbation problem for various types of (p, q) -generalized inverses in a Banach algebra.

Lemma 3.1 ([8, Lemma 2.2]). *Let $a, b \in \mathcal{A}$. If $1 + ab$ is left invertible, then so is $1 + ba$.*

Lemma 3.2. *Let $a, \delta a \in \mathcal{A}$ and $p, q \in \mathcal{A}^\bullet$ such that $a_{p,q}^{(2,l)}$ exists. Put $\bar{a} = a + \delta a$. If $1 + \delta a a_{p,q}^{(2,l)}$ is invertible, $w = a_{p,q}^{(2,l)}(1 + \delta a a_{p,q}^{(2,l)})^{-1}$. Then $\bar{a}_{p,q}^{(2,l)}$ exists and $w = \bar{a}_{p,q}^{(2,l)}$.*

Proof. We prove our result by showing that $waw = w$, $R_r(w) = R_r(p)$, $K_r(w) = R_r(q)$. It is easy to check that

$$w = a_{p,q}^{(2,l)}(1 + \delta a a_{p,q}^{(2,l)})^{-1} = (1 + a_{p,q}^{(2,l)}\delta a)^{-1} a_{p,q}^{(2,l)}.$$

Then, by using these two equalities, we can show $R_r(w) = R_r(a_{p,q}^{(2,l)}) = R_r(p)$ and $K_r(w) = K_r(a_{p,q}^{(2,l)}) = R_r(q)$. We can also compute

$$\begin{aligned}
 w\bar{a}w &= a_{p,q}^{(2,l)}(1 + \delta a a_{p,q}^{(2,l)})^{-1} \bar{a} a_{p,q}^{(2,l)}(1 + \delta a a_{p,q}^{(2,l)})^{-1} \\
 &= a_{p,q}^{(2,l)}(1 + \delta a a_{p,q}^{(2,l)})^{-1} [(a a_{p,q}^{(2,l)} - 1) + (1 + \delta a a_{p,q}^{(2,l)})] (1 + \delta a a_{p,q}^{(2,l)})^{-1} \\
 &= a_{p,q}^{(2,l)}(1 + \delta a a_{p,q}^{(2,l)})^{-1} (a a_{p,q}^{(2,l)} - 1) (1 + \delta a a_{p,q}^{(2,l)})^{-1} + w \\
 &= w.
 \end{aligned}$$

By Definition 1.1 and the uniqueness of $a_{p,q}^{(2,l)}$, we see $\bar{a}_{p,q}^{(2,l)}$ exists and $w = \bar{a}_{p,q}^{(2,l)}$. \square

Obviously, from the proof of Lemma 3.2, we see that if $a_{p,q}^{(2,l)}$ exists and $1 + a_{p,q}^{(2,l)}\delta a$ is invertible, set $v = (1 + a_{p,q}^{(2,l)}\delta a)^{-1} a_{p,q}^{(2,l)}$, then we also have $v = \bar{a}_{p,q}^{(2,l)}$. In order to prove the main results about the stable perturbation, we need one more characterizations of the existence of $a_{p,q}^{(2,l)}$.

For an element $a \in \mathcal{A}$ and $p, q \in \mathcal{A}^\bullet$. Let $R_a : \mathcal{A} \rightarrow \mathcal{A}$ be the right multiplier on \mathcal{A} (i.e., $R_a(x) = xa$ for any $x \in \mathcal{A}$). Then, it easy to see that $a_{p,q}^{(2,l)}$ exists in \mathcal{A} if and only if $(R_a)_{\mathcal{A}(1-q), \mathcal{A}(1-p)}^{(2)}$ exists in the Banach algebra $B(\mathcal{A})$. So from the equivalences of (1), (2), and (3) in Lemma 2.2, dually, we can get the following equivalent conditions for the existence of $a_{p,q}^{(2,l)}$.

Proposition 3.3. *Let $a \in \mathcal{A}$ and $p, q \in \mathcal{A}^\bullet$. Then the following statements are equivalent:*

(1) $a_{p,q}^{(2,l)}$ exists;

(2) there exists $c \in \mathcal{A}$ such that $cac = c$, $R_l(c) = R_l(1 - q)$, and $K_l(c) = R_l(1 - p)$;

(3) $K_l(a) \cap R_l(1 - q) = \{0\}$ and $\mathcal{A} = R_l(1 - q)a + R_l(1 - p)$.

Proof. (1) \Leftrightarrow (2) Suppose that $a_{p,q}^{(2,l)}$ exists. Let $c = a_{p,q}^{(2,l)}$. Then from Definition 1.1, we know that $cac = c$, and then $ca, ac \in \mathcal{A}^\bullet$, $R_r(ca) = R_r(c) = R_r(p)$, $K_r(ac) = K_r(c) = R_r(q)$. Thus, it follows from Lemma 2.1 that

$$cap = p, \quad pca = ca, \quad ac(1 - q) = ac, \quad (1 - q)ac = 1 - q.$$

Then, by using Lemma 2.1 again, we have

$$K_l(ca) \subset K_l(p) \subset K_l(ca), \quad R_l(ac) \subset R_l(1 - q) \subset R_l(ac). \quad (3.1)$$

By using $cac = c$, we have $K_l(ca) = K_l(c)$ and $R_r(ac) = R_r(c)$. Thus from Equation (3.1), we see that (2) holds. If (2) holds, similarly, by using Definition 1.1 and Lemma 2.1, we can obtain $a_{p,q}^{(2,l)}$ exists.

(2) \Leftrightarrow (3) By our remark above this lemma, we see these hold simply from the equivalences of (2) and (3) in Lemma 2.2. Note that we can also prove these equivalences directly by using the right multiplier R_a on \mathcal{A} . Here we omit the detail. \square

Now, we can present one of our main results about the stable perturbation of the generalized inverse $a_{p,q}^{(2,l)}$.

Theorem 3.4. Let $a, \delta a \in \mathcal{A}$ and $p, q \in \mathcal{A}^\bullet$ such that $a_{p,q}^{(2,l)}$ exists. Put $\bar{a} = a + \delta a$. Then the following statements are equivalent:

(1) $1 + \delta a a_{p,q}^{(2,l)}$ is invertible;

(2) $1 + a_{p,q}^{(2,l)}\delta a$ is invertible;

(3) $\bar{a}_{p,q}^{(2,l)}$ exists.

In this case, we have $\bar{a}_{p,q}^{(2,l)} = a_{p,q}^{(2,l)}(1 + \delta a a_{p,q}^{(2,l)})^{-1} = (1 + a_{p,q}^{(2,l)}\delta a)^{-1} a_{p,q}^{(2,l)}$.

Proof. (1) \Leftrightarrow (2) Follows from the well-known spectral theory in Banach algebras.

(2) \Rightarrow (3) We prove our result by using Lemma 2.2. Let $x \in K_r(\bar{a}) \cap R_r(p) = \{0\}$. Since $R_r(p) = R_r(a_{p,q}^{(2,l)})$, then there exists some $t \in \mathcal{A}$ satisfying $x = a_{p,q}^{(2,l)}t$ and $\bar{a}t = 0$. Thus, we have

$$\begin{aligned} (1 + a_{p,q}^{(2,l)}\delta a)a_{p,q}^{(2,l)}t &= a_{p,q}^{(2,l)}t + a_{p,q}^{(2,l)}\delta a a_{p,q}^{(2,l)}t \\ &= a_{p,q}^{(2,l)}(a + \delta a)a_{p,q}^{(2,l)}t \\ &= a\bar{a}t = 0. \end{aligned}$$

Since $1 + a_{p,q}^{(2,l)}\delta a$ is invertible, it follows that $x = a_{p,q}^{(2,l)}t = 0$. Therefore,

$$K_r(\bar{a}) \cap R_r(p) = \{0\}. \quad (3.2)$$

Let $s \in \bar{a}R_r(p) \cap R_r(q)$. Since $R_r(p) = R_r(a_{p,q}^{(2,l)})$ and $R_r(q) = K_r(a_{p,q}^{(2,l)})$, then there exists some $z \in \mathcal{A}$ such that $s = \bar{a}a_{p,q}^{(2,l)}z$ and $a_{p,q}^{(2,l)}s = 0$.

Similar to the proof of Equation (3.2), we can get $s = a_{p,q}^{(2,l)}t = 0$, i.e.,

$\bar{a}R_r(p) \cap R_r(q) = \{0\}$. Since $1 + a_{p,q}^{(2,l)}\delta a$ is invertible, then for any

$w \in \mathcal{A}$, there is some $v \in \mathcal{A}$ such that $a_{p,q}^{(2,l)}w = (1 + a_{p,q}^{(2,l)}\delta a)v$. From $\bar{a} = a + \delta a$, we have

$$(1 - a_{p,q}^{(2,l)}a)v = a_{p,q}^{(2,l)}(w - \bar{a}v) \in R_r(a_{p,q}^{(2,l)}a) \cap K_r(a_{p,q}^{(2,l)}a) = \{0\}.$$

Thus, $w - \bar{a}v \in K_r(a_{p,q}^{(2,l)})$ and $v = a_{p,q}^{(2,l)}av \in R_r(a_{p,q}^{(2,l)})$. Since for $w \in \mathcal{A}$, we also have $w = \bar{a}v + (w - \bar{a}v) \in \bar{a}R_r(p) \dot{+} R_r(q)$. Thus, we have

$$\mathcal{A} = \bar{a}R_r(p) \dot{+} R_r(q). \quad (3.3)$$

Now, from Equations (3.2) and (3.3), by using Lemma 2.2, we see that $\bar{a}_{p,q}^{(2,l)}$ exists.

(3) \Rightarrow (1) Suppose that $\bar{a}_{p,q}^{(2,l)}$ exists, we want to prove $1 + \delta a a_{p,q}^{(2,l)}$ is both left and right invertible. Since $\bar{a}_{p,q}^{(2,l)}$ exists, then from Lemma 2.2, $\mathcal{A} = \bar{a}R_r(p) \dot{+} R_r(q) = \bar{a}R_r(a_{p,q}^{(2,l)}) \dot{+} K_r(a_{p,q}^{(2,l)})$. Thus, for any $x \in \mathcal{A}$, we can write $x = \bar{a}a_{p,q}^{(2,l)}t_1 + t_2$, where $t_1 \in \mathcal{A}$ and $t_2 \in K_r(a_{p,q}^{(2,l)})$. Set $s = a a_{p,q}^{(2,l)}t_1 + t_2$, then

$$\begin{aligned} (1 + \delta a a_{p,q}^{(2,l)})s &= (1 + \delta a a_{p,q}^{(2,l)})(a a_{p,q}^{(2,l)}t_1 + t_2) \\ &= \bar{a}a_{p,q}^{(2,l)}t_1 + t_2 = x. \end{aligned}$$

Since $x \in \mathcal{A}$ is arbitrary, let $x = 1$, then we see that $1 + \delta a a_{p,q}^{(2,l)}$ is right invertible. Now we prove that $1 + \delta a a_{p,q}^{(2,l)}$ is also left invertible. In fact, from Proposition 3.3, we also have $\mathcal{A} = R_l(1 - q)\bar{a} \dot{+} R_l(1 - p) = R_l(a_{p,q}^{(2,l)})\bar{a} \dot{+} K_l(a_{p,q}^{(2,l)})$ for $\bar{a}_{p,q}^{(2,l)}$ exists. Then for any $z \in \mathcal{A}$, we can write $z = s_1 a_{p,q}^{(2,l)}\bar{a} + s_2$, where $s_1 \in R_l(a_{p,q}^{(2,l)}a)$ and $s_2 \in K_l(a_{p,q}^{(2,l)})$. Let $t = s_1 + s_2$, then we have

$$\begin{aligned} t(1 + a_{p,q}^{(2,l)}\delta a) &= (s_1 + s_2)(1 + a_{p,q}^{(2,l)}\delta a) \\ &= s_1 + s_2 + s_1 a_{p,q}^{(2,l)}(\bar{a} - a) \\ &= s_1 a_{p,q}^{(2,l)}\bar{a} + s_2 + s_1(1 - a_{p,q}^{(2,l)}a) \\ &= z. \end{aligned}$$

Since $z \in \mathcal{A}$ is arbitrary, let $z = 1$, then we get that $1 + a_{p,q}^{(2,l)}\delta a$ is left invertible. But from Lemma 3.1, we see $1 + \delta a a_{p,q}^{(2,l)}$ is also left invertible.

Thus, $1 + \delta a a_{p,q}^{(2,l)}$ is invertible.

Now, from Lemma 3.2, $\bar{a}_{p,q}^{(2,l)} = a_{p,q}^{(2,l)}(1 + \delta a a_{p,q}^{(2,l)})^{-1} = (1 + a_{p,q}^{(2,l)}\delta a)^{-1} a_{p,q}^{(2,l)}$. This completes the proof. \square

Lemma 3.5. *Let $a, \delta a \in \mathcal{A}$ and $p, q \in \mathcal{A}^\bullet$ such that $a_{p,q}^{(2,l)}$ exists. If $1 + a_{p,q}^{(2,l)}\delta a$ is invertible. Put $\bar{a} = a + \delta a$ and $f = (1 + a_{p,q}^{(2,l)}\delta a)^{-1}(1 - a_{p,q}^{(2,l)}a)$. Then*

- (1) $f \in \mathcal{A}^\bullet$ with $K_r(\bar{a}) \subset R_r(f)$;
- (2) $K_r(\bar{a}) = R_r(f)$ if and only if $R_r(\bar{a}) \cap R_r(q) = \{0\}$.

Proof. (1) Since $(1 - a_{p,q}^{(2,l)}a)(1 + a_{p,q}^{(2,l)}\delta a) = 1 - a_{p,q}^{(2,l)}a$ and $1 + a_{p,q}^{(2,l)}\delta a$ is invertible, we have $1 - a_{p,q}^{(2,l)}a = (1 - a_{p,q}^{(2,l)}a)(1 + a_{p,q}^{(2,l)}\delta a)^{-1}$. Thus,

$$f^2 = (1 + a_{p,q}^{(2,l)}\delta a)^{-1}(1 - a_{p,q}^{(2,l)}a)(1 + a_{p,q}^{(2,l)}\delta a)^{-1}(1 - a_{p,q}^{(2,l)}a) = f.$$

Now for any $x \in \mathcal{A}$, from $(1 - a_{p,q}^{(2,l)}a)x = (1 + a_{p,q}^{(2,l)}\delta a)x - a_{p,q}^{(2,l)}\bar{a}x$, we have

$$fx = (1 + a_{p,q}^{(2,l)}\delta a)^{-1}(1 - a_{p,q}^{(2,l)}a)x = x - (1 + a_{p,q}^{(2,l)}\delta a)^{-1}a_{p,q}^{(2,l)}\bar{a}x. \quad (3.4)$$

Equation (3.4) implies that $K_r(\bar{a}) \subset R_r(f)$.

(2) (\Rightarrow) Let $t \in R_r(\bar{a}) \cap R_r(q) = \{0\}$. Since $R_r(q) = K_r(a_{p,q}^{(2,l)})$, then there is some $x \in \mathcal{A}$ such that $t = \bar{a}x$ and $a_{p,q}^{(2,l)}\bar{a}x = a_{p,q}^{(2,l)}t = 0$. Thus, $x = fx$ by Equation (3.4). So, $x \in R_r(f) = K_r(\bar{a})$ and $t = \bar{a}x = 0$, i.e., $R_r(\bar{a}) \cap R_r(q) = \{0\}$.

(\Leftarrow) Thanks to (1), we need only to prove $R_r(f) \subset K_r(\bar{a})$. Let $t \in R_r(f)$. Since $f \in \mathcal{A}^\bullet$, we have $t = ft$. So by Equation (3.4), we get $(1 + a_{p,q}^{(2,l)}\delta a)^{-1} a_{p,q}^{(2,l)} \bar{a} t = 0$ and then $a_{p,q}^{(2,l)} \bar{a} t = 0$. Hence $\bar{a} t \in R_r(\bar{a}) \cap R_r(q) = \{0\}$ and $K_r(\bar{a}) = R_r(f)$. \square

Similar to [12, Proposition 2.2] or [13, Theorem 2.4.7], but by using some of our characterizations for $a_{p,q}^{(2,l)}$ and $a_{p,q}^{(l)}$, we can obtain the following results about the stable perturbations for these two kinds of generalized inverses.

Theorem 3.6. *Let $a, \delta a \in \mathcal{A}$ and $p, q \in \mathcal{A}^\bullet$ such that $a_{p,q}^{(2,l)}$ exists. Suppose that $1 + a_{p,q}^{(2,l)}\delta a$ is invertible. Put $\bar{a} = a + \delta a$ and $w = (1 + a_{p,q}^{(2,l)}\delta a)^{-1} a_{p,q}^{(2,l)}$. Then the following statements are equivalent:*

- (1) $w = \bar{a}_{p,q}^{(l)}$, i.e., $\bar{a}_{p,q}^{(2,l)} = \bar{a}_{p,q}^{(l)}$;
- (2) $R_r(\bar{a}) \cap R_r(q) = \{0\}$, i.e., \bar{a} is a stable perturbation of a ;
- (3) $\bar{a}(1 + a_{p,q}^{(2,l)}\delta a)^{-1}(1 - a_{p,q}^{(2,l)}a) = 0$;
- (4) $(1 - aa_{p,q}^{(2,l)})(1 + \delta aa_{p,q}^{(2,l)})^{-1}\bar{a} = 0$.

Proof. The implication (1) \Leftrightarrow (2) comes from Lemmas 2.2, 2.3, and Theorem 3.4. The implication (2) \Leftrightarrow (3) comes from Lemma 3.5.

(3) \Leftrightarrow (4) We can compute in the following way:

$$\begin{aligned}
\bar{a}(1 + a_{p,q}^{(2,l)}\delta a)^{-1}(1 - a_{p,q}^{(2,l)}a) &= \bar{a}(1 + a_{p,q}^{(2,l)}\delta a)^{-1}[(1 + a_{p,q}^{(2,l)}\delta a) - a_{p,q}^{(2,l)}\bar{a}] \\
&= \bar{a} - \bar{a}a_{p,q}^{(2,l)}(1 + \delta aa_{p,q}^{(2,l)})^{-1}\bar{a} \\
&= \bar{a} - [(1 + \delta aa_{p,q}^{(2,l)}) - (1 - aa_{p,q}^{(2,l)})]
\end{aligned}$$

$$\begin{aligned}
 & \times (1 + \delta a a_{p,q}^{(2,l)})^{-1} \bar{a} \\
 & = (1 - a a_{p,q}^{(2,l)}) (1 + \delta a a_{p,q}^{(2,l)})^{-1} \bar{a}.
 \end{aligned}$$

This completes the proof. \square

Furthermore, by using the above theorem, we have the following results:

Corollary 3.7. *Let $a, \delta a \in \mathcal{A}$ and $p, q \in \mathcal{A}^\bullet$ such that $a_{p,q}^{(2,l)}$ exists. Put $\bar{a} = a + \delta a$. If $1 + a_{p,q}^{(2,l)} \delta a$ is invertible. Then the following statements are equivalent:*

- (1) $R_r(\bar{a}) \cap R_r(q) = \{0\}$, i.e., \bar{a} is stable perturbation of a ;
- (2) $(1 + a_{p,q}^{(2,l)} \delta a)^{-1} K_r(a_{p,q}^{(2,l)} a) = K_r(\bar{a})$;
- (3) $(1 + \delta a a_{p,q}^{(2,l)})^{-1} R_r(\bar{a}) = R_r(a a_{p,q}^{(2,l)})$.

Proof. Note that we have $K_r(a_{p,q}^{(2,l)} a) = R_r(1 - a_{p,q}^{(2,l)} a)$ and $R_r(a a_{p,q}^{(2,l)}) = K_r(1 - a a_{p,q}^{(2,l)})$. So, we can get the assertions by using Theorem 3.6. \square

Theorem 3.8. *Let $a, \delta a \in \mathcal{A}$ and $p, q \in \mathcal{A}^\bullet$ such that $a_{p,q}^{(l)}$ exists. Put $\bar{a} = a + \delta a$. Then the following statements are equivalent:*

- (1) $1 + a_{p,q}^{(l)} \delta a$ is invertible, $R_r(\bar{a}) = K_r(q)$ and $\bar{a}_{p,q}^{(l)} = a_{p,q}^{(l)} (1 + \delta a a_{p,q}^{(l)})^{-1}$.
- (2) $R_r(\bar{a}) \cap R_r(q) = \{0\}$, $K_r(\bar{a}) \cap R_r(p) = \{0\}$, and $\bar{a} R_r(p) = K_r(q)$.

Proof. (1) \Rightarrow (2) Suppose that (1) holds. Since $a_{p,q}^{(l)}$ exists, we obtain that $a_{p,q}^{(2,l)}$ exists and $a_{p,q}^{(2,l)} = a_{p,q}^{(l)}$. Thus, from our assumption, by using Theorem 3.6 and Lemma 2.3, we have

$$R_r(\bar{a}) \cap R_r(q) = \{0\}, \quad K_r(\bar{a}) \cap R_r(p) = \{0\}.$$

Now we need to show $\bar{a}R_r(p) = K_r(q)$. But since $R_r(\bar{a}) = K_r(q)$, so we can prove our result by showing that $\bar{a}R_r(p) = R_r(\bar{a})$. Obviously, $\bar{a}R_r(p) \subset R_r(\bar{a})$. On the other hand, since $\bar{a}_{p,q}^{(l)}$ exists, then by Lemma 2.3 again, we have $\mathcal{A} = \bar{a}R_r(p) \dot{+} R_r(q)$. Now for any $x \in R_r(\bar{a})$, we can write $x = x_1 + x_2$ with $x_1 \in \bar{a}R_r(p)$ and $x_2 \in R_r(q)$. From $\bar{a}R_r(p) \subset R_r(\bar{a})$, we get $x_1 \in R_r(\bar{a})$. Thus,

$$x_2 = x - x_1 \in R_r(\bar{a}) \cap R_r(q) = \{0\}.$$

Therefore, $x_2 = 0$ and then $x = x_1 \in \bar{a}R_r(p)$. Hence, $\bar{a}R_r(p) = R_r(\bar{a}) = K_r(q)$.

(2) \Rightarrow (1) Since $q \in \mathcal{A}^\bullet$ and $\bar{a}R_r(p) = K_r(q)$, we can write $\mathcal{A} = K_r(q) \dot{+} R_r(q) = \bar{a}R_r(p) \dot{+} R_r(q)$. Note that $R_r(\bar{a}) \cap R_r(q) = \{0\}$, $K_r(\bar{a}) \cap R_r(p) = \{0\}$, then by using Lemma 2.3, we get $a_{p,q}^{(l)}$ exists, then $a_{p,q}^{(2,l)}$ also exists and $a_{p,q}^{(2,l)} = a_{p,q}^{(l)}$. Thus, from Theorem 3.4, we see $1 + a_{p,q}^{(l)}\delta a$ is invertible and $\bar{a}_{p,q}^{(l)} = a_{p,q}^{(l)}(1 + \delta a a_{p,q}^{(l)})^{-1}$. Now, by using Lemma 2.3, we can get $\mathcal{A} = \bar{a}R_r(p) \dot{+} R_r(q)$. Similarly, as in (1) \Rightarrow (2), by using $R_r(\bar{a}) \cap R_r(q) = \{0\}$, we can show that $\bar{a}R_r(p) = R_r(\bar{a})$ and then $R_r(\bar{a}) = K_r(q)$. This completes the proof. \square

The first result in the following lemma has been proved for generalized inverse a^+ by the second author (see [12, Proposition 2.5]). By using the same method, we can prove the following results for the general inverse $a_{p,q}^{(l)}$.

Lemma 3.9 ([12, Proposition 2.5]). *Let $a, \delta a \in \mathcal{A}$ and $p, q \in \mathcal{A}^\bullet$ such that $a_{p,q}^{(l)}$ exists. Put $\bar{a} = a + \delta a$.*

(1) If $\delta(R_r(\bar{a}), R_r(a)) < \|1 - aa_{p,q}^{(l)}\|^{-1}$, then $R_r(\bar{a}) \cap R_r(q) = \{0\}$;

(2) If $\delta(K_r(\bar{a}), K_r(a)) < \|a_{p,q}^{(l)}a\|^{-1}$, then $K_r(\bar{a}) \cap R_r(p) = \{0\}$.

By using Theorems 3.6, 3.8, and Lemma 3.9, we have the following:

Corollary 3.10. Let $a, \delta a \in \mathcal{A}$ and $p, q \in \mathcal{A}^\bullet$ such that $a_{p,q}^{(l)}$ exists.

Put $\bar{a} = a + \delta a$. If one of the following condition holds:

(i) $\delta(K_r(\bar{a}), K_r(a)) < \|a_{p,q}^{(l)}a\|^{-1}$, $\delta(R_r(\bar{a}), R_r(a)) < \|1 - aa_{p,q}^{(l)}\|^{-1}$, and $\bar{a}R_r(p) = K_r(q)$.

(ii) $1 + a_{p,q}^{(l)}\delta a$ is invertible and $\delta(R_r(\bar{a}), R_r(a)) < \|1 - aa_{p,q}^{(l)}\|^{-1}$.

Then $\bar{a}_{p,q}^{(l)}$ exists and $\bar{a}_{p,q}^{(l)} = a_{p,q}^{(l)}(1 + \delta a a_{p,q}^{(l)})^{-1}$.

Finally, we present some perturbation results for $a_{p,q}^{(2)}$.

Theorem 3.11. Let $a, \delta a \in \mathcal{A}$ and $p, q \in \mathcal{A}^\bullet$ such that $a_{p,q}^{(2)}$ exists.

Put $\bar{a} = a + \delta a$. If $1 + a_{p,q}^{(2)}\delta a$ is invertible. Then the following statements are equivalent:

(1) $\bar{a}_{p,q}^{(2)}$ exists and $\bar{a}_{p,q}^{(2)} = a_{p,q}^{(2)}(1 + \delta a a_{p,q}^{(2)})^{-1}$;

(2) $\bar{a}p = (1 - q)\bar{a}$;

(3) $\bar{a}a_{p,q}^{(2)} = (1 - q)\bar{a}a_{p,q}^{(2)}$ and $a_{p,q}^{(2)}\bar{a} = a_{p,q}^{(2)}\bar{a}p$.

Proof. (1) \Leftrightarrow (2) Comes from [6, Theorem 4.1]. We show that (2) and (3) are equivalent. If $\bar{a}p = (1 - q)\bar{a}$, then

$$\bar{a}a_{p,q}^{(2)} = \bar{a}p a_{p,q}^{(2)} = (1 - q)\bar{a}a_{p,q}^{(2)} \quad \text{and} \quad a_{p,q}^{(2)}\bar{a} = a_{p,q}^{(2)}(1 - q)\bar{a} = a_{p,q}^{(2)}\bar{a}p.$$

Conversely, if (3) holds, then $\bar{a}p = \bar{a}a_{p,q}^{(2)}a = (1-q)\bar{a}a_{p,q}^{(2)}a = aa_{p,q}^{(2)}\bar{a}p = (1-q)\bar{a}$. \square

Corollary 3.12. *Let $a, \delta a \in \mathcal{A}$ and $p, q \in \mathcal{A}^\bullet$ such that $a_{p,q}^{(2)}$ exists. Put $\bar{a} = a + \delta a$. If $1 + a_{p,q}^{(2)}\delta a$ is invertible and $\delta a = (1-q)\delta a = \delta ap$. Then $\bar{a}_{p,q}^{(2)}$ exists and*

$$\bar{a}_{p,q}^{(2)} = a_{p,q}^{(2)}(1 + \delta a a_{p,q}^{(2)})^{-1} = (1 + a_{p,q}^{(2)}\delta a)^{-1}a_{p,q}^{(2)}.$$

Proof. If $\delta a = (1-q)\delta a = \delta ap$, then it is easy to check that $\bar{a}p = (1-q)\bar{a}$. Thus, Theorem 3.11 shows that our results hold. \square

4. Perturbation Analysis for the (p, q) -Generalized Inverses

In this section, we mainly investigate the general perturbations problem for the (p, q) -generalized inverses $a_{p,q}^{(2,l)}$ and $a_{p,q}^{(1,2)}$. Let $\kappa = \|a\| \|a_{p,q}^{(2,l)}\|$, which is the generalized condition number of the generalized inverse $a_{p,q}^{(2,l)}$.

Lemma 4.1. *Let $a \in \mathcal{A}$ and $p \in \mathcal{A}^\bullet$ with $R_r(p) = R_r(a)$. Let $c \in \mathcal{A}$ with $R_r(c)$ closed and $\hat{\delta}(R_r(c), R_r(a)) < \frac{1}{1 + \|p\|}$. Then $\mathcal{A} = R_r(c) \dot{+} K_r(p)$.*

Proof. Let $L_p x = px, \forall x \in \mathcal{A}$. Then L_p is an idempotent operator on \mathcal{A} with $\|L_p\| = \|p\|$ and $\{L_p x \mid x \in \mathcal{A}\} = R_r(p)$. By [10, Theorem 11] or [13, Lemma 4.4.4], $\hat{\delta}(R_r(c), R_r(a)) < \frac{1}{1 + \|L_p\|}$ implies that $\mathcal{A} = R_r(c) \dot{+} K_r(p)$. \square

Lemma 4.2 ([12, Lemma 2.4]). *For any $p, q \in \mathcal{A}^\bullet$, we have $\hat{\delta}(R_r(p), R_r(q)) \leq \|p - q\|$.*

Lemma 4.3. *Let $a \in \mathcal{A}$ and $p, q \in \mathcal{A}^\bullet$ such that $\alpha_{p,q}^{(2,l)}$ exists.*

Suppose that $p' \in \mathcal{A}^\bullet$ satisfying $\|p - p'\| < \frac{1}{1 + \kappa}$. Then

$$(1) \hat{\delta}(aR_r(p), aR_r(p')) \leq \frac{\kappa\|p - p'\|}{1 - (1 + \kappa)\|p - p'\|};$$

$$(2) aR_r(p') \subset \mathcal{A} \text{ is closed and } K_r(a) \cap R_r(p') = \{0\}.$$

Proof. (1) Set $b = \alpha_{p,q}^{(2,l)}$. For any $t' \in R_r(p')$, we have

$$\begin{aligned} \text{dist}(at', aR_r(p)) &= \inf_{t \in R_r(p)} \|at' - at\| \leq \|a\| \inf_{t \in R_r(p)} \|t' - t\| \\ &\leq \|a\| \text{dist}(t', R_r(p)) \\ &\leq \|a\| \|t'\| \hat{\delta}(R_r(p'), R_r(p)). \end{aligned}$$

Thus, we get

$$\delta(at', aR_r(p)) \leq \|a\| \|t'\| \hat{\delta}(R_r(p'), R_r(p)). \quad (4.1)$$

But for any $t' \in R_r(p')$ and $t \in R_r(p)$, we have

$$\begin{aligned} \|b\| \|at'\| &= \|b\| \|a(t' - t + t)\| \geq \|b\| \|at\| - \|b\| \|a\| \|t' - t\| \\ &\geq \|t\| - \|b\| \|a\| \|t' - t\| \\ &\geq \|t'\| - (1 + \|b\| \|a\|) \|t' - t\|. \end{aligned}$$

Thus, $\|t'\| - \|b\| \|at'\| \leq (1 + \|b\| \|a\|) \|t' - t\|$, and then

$$\begin{aligned} \|t'\| - \|b\| \|at'\| &\leq (1 + \|b\| \|a\|) \text{dist}(t', R_r(p)) \\ &\leq (1 + \|b\| \|a\|) \|t'\| \hat{\delta}(R_r(p'), R_r(p)). \end{aligned}$$

Therefore, we have

$$\|t'\| \leq \frac{\|b\| \|at'\|}{1 - (1 + \|b\| \|a\|) \hat{\delta}(R_r(p'), R_r(p))}. \quad (4.2)$$

Then by Equations (4.1) and (4.2), we get

$$\delta(at', aR_r(p)) \leq \frac{\|b\| \|a\| \|at'\| \hat{\delta}(R_r(p'), R_r(p))}{1 - (1 + \|b\| \|a\|) \hat{\delta}(R_r(p'), R_r(p))}.$$

Now, by Lemma 4.2 and the definition of gap-function, we have

$$\delta(aR_r(p'), aR_r(p)) \leq \frac{\kappa \|p - p'\|}{1 - (1 + \kappa) \|p - p'\|}. \quad (4.3)$$

On the other hand, for any $t \in R_r(p)$, by Lemma 2.2, we have $t = pt = bap t = bat$, then

$$\begin{aligned} \text{dist}(at, aR_r(p')) &= \inf_{s \in R_r(p')} \|at - as\| \leq \|a\| \inf_{s \in R_r(p')} \|t - s\| \\ &= \|a\| \text{dist}(t, R_r(p')) \leq \|a\| \|t\| \delta(R_r(p), R_r(p')) \\ &= \|a\| \|bat\| \delta(R_r(p), R_r(p')) \\ &\leq \|a\| \|b\| \|at\| \delta(R_r(p), R_r(p')). \end{aligned}$$

Thus, we have $\delta(aR_r(p), aR_r(p')) \leq \kappa \hat{\delta}(R_r(p), R_r(p'))$. So by Lemma 4.2,

$$\delta(aR_r(p), aR_r(p')) \leq \kappa \|p - p'\|. \quad (4.4)$$

Consequently, from Equations (4.3) and (4.4), we have

$$\begin{aligned} \hat{\delta}(aR_r(p), aR_r(p')) &= \max \{ \delta(aR_r(p'), aR_r(p)), \delta(aR_r(p), aR_r(p')) \} \\ &\leq \frac{\kappa \|p - p'\|}{1 - (1 + \kappa) \|p - p'\|}. \end{aligned}$$

(2) Obviously, by Equation (4.2), we get $aR_r(p') \subset \mathcal{A}$ is closed and $K_r(a) \cap R_r(p') = \{0\}$. This completes the proof. \square

Now, we can give the following perturbation result for $a_{p,q}^{(2,l)}$ when p has a small perturbation.

Theorem 4.4. *Let $a \in \mathcal{A}$ and $p, q \in \mathcal{A}^\bullet$ such that $a_{p,q}^{(2,l)}$ exists.*

Suppose that $p' \in \mathcal{A}^\bullet$ with $\|p - p'\| < \frac{1}{(1 + \kappa)^2}$. Then $a_{p',q}^{(2,l)}$ exists and

$$\frac{\|a_{p',q}^{(2,l)} - a_{p,q}^{(2,l)}\|}{\|a_{p,q}^{(2,l)}\|} \leq \frac{(1 + \kappa)\|p' - p\|}{1 - (1 + \kappa)\|p' - p\|} \text{ and } \|a_{p',q}^{(2,l)}\| \leq \frac{\|a_{p,q}^{(2,l)}\|}{1 - (1 + \kappa)\|p' - p\|}.$$

Proof. Let $b = a_{p,q}^{(2,l)}$, then by Definition 1.1, we know that $ab \in \mathcal{A}^\bullet$ and $\kappa = \|b\|\|a\| \geq \|ab\|$. By using Lemma 4.3 and note that $\|p - p'\| < \frac{1}{(1 + \kappa)^2}$, we have

$$\hat{\delta}(aR_r(p), aR_r(p')) \leq \frac{\kappa\|p - p'\|}{1 - (1 + \kappa)\|p - p'\|} < \frac{1}{1 + \|ab\|}.$$

From Lemma 4.3 (2), we know that $K_r(a) \cap R_r(p') = \{0\}$ and $aR_r(p') \subset \mathcal{A}$ is closed. Thus, by Lemma 4.1, $aR_r(p')$ is complemented and $\mathcal{A} = aR_r(p') \dot{+} R_r(q)$. Therefore, by Lemma 2.2, we know $a_{p',q}^{(2)}$ exists.

For convenience, we write $a_{p',q}^{(2,l)} = b'$. Since we have proved that $a_{p',q}^{(2,l)}$ exists, then by Theorem 2.2, $\mathcal{A} = aR_r(p') \dot{+} R_r(q) = aR_r(p') \dot{+} K_r(b')$ and $K_r(b') = R_r(q) = K_r(b)$. Thus for any $x \in \mathcal{A}$, we can write $x = t + t'$ with $t = ab'z$ for some $z \in \mathcal{A}$ and $t' \in R_r(q)$.

Since $\text{dist}(b'z, R_r(p)) \leq \|b'z\|\delta(R_r(p'), R_r(p)) \leq \|b'z\|\|p' - p\|$, then for every $\epsilon > 0$, we can choose $y \in \mathcal{A}$ such that $\|b'z - by\| < \|b'z\|\|p' - p\| + \epsilon$. Put $s = aby$. Then, we have

$$\|t - s\| = \|ab'z - aby\| < \|a\|\|b'z\|\|p' - p\| + \|a\|\epsilon.$$

$$\begin{aligned} \|(b' - b)x\| &= \|(b' - b)(t + t')\| = \|(b' - b)t\| \\ &\leq \|b'ab't - babs\| + \|bs - bt\| \end{aligned}$$

$$\begin{aligned}
&\leq \|b'z - by\| + \|b\| \|s - t\| \\
&\leq (1 + \kappa) \|b'z\| \|p' - p\| + (1 + \kappa)\epsilon.
\end{aligned} \tag{4.5}$$

From $t = ab'z$, we get $b't = b'z$ and therefore,

$$\|b'z\| = \|b't\| = \|(b' - b)x + bx\| \leq \|(b' - b)x\| + \|b\| \|x\|. \tag{4.6}$$

Thus by using Equations (4.5) and (4.6), we get

$$\|(b' - b)x\| \leq (1 + \kappa)(\|b\| \|x\| + \|(b' - b)x\|) \|p' - p\| + (1 + \kappa)\epsilon. \tag{4.7}$$

Letting $\epsilon \rightarrow 0^+$ in Equation (4.7), we can get

$$\frac{\|a_{p',q}^{(2,l)} - a_{p,q}^{(2,l)}\|}{\|a_{p,q}^{(2,l)}\|} \leq \frac{(1 + \kappa) \|p' - p\|}{1 - (1 + \kappa) \|p' - p\|}, \quad \|a_{p',q}^{(2,l)}\| \leq \frac{\|a_{p,q}^{(2,l)}\|}{1 - (1 + \kappa) \|p' - p\|}.$$

This completes the proof. \square

Some representations for the generalized inverse $\alpha_{p,q}^{(2,l)}$ have been presented in [1]. The following result gives a representation of $\alpha_{p,q}^{(2,l)}$ based on (1, 5) inverse. Note that this result is also an improvement of the group inverses representation of $\alpha_{p,q}^{(2,l)}$ (see [1]), which removes the existence of the group inverses of wa or aw .

Lemma 4.5 ([1, Theorem 5.6]). *Let $a, w \in \mathcal{A}$ and $p, q \in \mathcal{A}^\bullet$ such that $R_r(w) = R_r(p)$ and $K_r(w) = R_r(q)$. Then the following statements are equivalent:*

- (1) $\alpha_{p,q}^{(2,l)}$ exists;
- (2) $(aw)^{(1,5)}$ exists and $K_r(a) \cap R_r(w) = \{0\}$;
- (3) $(wa)^{(1,5)}$ exists and $R_r(w) = R_r(wa)$.

In this case, waw is inner regular and

$$\alpha_{p,q}^{(2,l)} = (wa)^{(1,5)}w = w(aw)^{(1,5)} = w(waw)^-w.$$

Now, we give the result when q has a small perturbation. By using our above Lemma 4.5, we can also give a new representation for the generalized inverse of the perturbed operator.

Theorem 4.6. *Let $a \in \mathcal{A}$ and $p, q \in \mathcal{A}^\bullet$ such that $\alpha_{p,q}^{(2,l)}$ exists.*

Suppose that $q' \in \mathcal{A}^\bullet$ with $\|q - q'\| < \frac{1}{2 + \kappa}$. Then $\alpha_{p,q'}^{(2,l)}$ exists and

$$(1) \frac{\|\alpha_{p,q'}^{(2,l)} - \alpha_{p,q}^{(2,l)}\|}{\|\alpha_{p,q}^{(2,l)}\|} \leq \frac{(1 + \kappa)\|q - q'\|}{1 - \kappa\|q - q'\|} \text{ and } \|\alpha_{p,q'}^{(2,l)}\| \leq \frac{1 + \|q' - q\|}{1 - \kappa\|q' - q\|} \|\alpha_{p,q}^{(2,l)}\|.$$

(2) *If there are some $w, v \in \mathcal{A}$ with $R_r(w) = R_r(p)$, $K_r(w) = K_r(v) = R_r(q)$ and $R_r(v) = R_r(q')$. Then*

$$\alpha_{p,q'}^{(2,l)} = \alpha_{p,q}^{(2,l)} + \alpha_{p,q}^{(2,l)}(av)^{(1,5)}a(v - w)(1 - a\alpha_{p,q}^{(2,l)}).$$

Proof. Since $1 + \|1 - ab\| \leq 2 + \kappa$, then by Lemma 4.2, we have

$$\hat{\delta}(R_r(q), R_r(q')) \leq \|q - q'\| < \frac{1}{2 + \kappa} \leq \frac{1}{1 + \|1 - ab\|}.$$

So $\mathcal{A} = aR_r(p) \dot{+} R_r(q')$ by Lemma 4.1. Note that $K_r(a) \cap R_r(p) = \{0\}$.

So $\alpha_{p,q'}^{(1,2)}$ exists.

(1) From $\alpha_{p,q'}^{(2,l)} = p\alpha_{p,q'}^{(2,l)} = \alpha_{p,q}^{(2,l)}a\alpha_{p,q'}^{(2,l)}$, we get

$$\alpha_{p,q'}^{(2,l)} - \alpha_{p,q}^{(2,l)} = \alpha_{p,q}^{(2,l)}(a\alpha_{p,q'}^{(2,l)} - a\alpha_{p,q}^{(2,l)}).$$

Since $(1 - a\alpha_{p,q'}^{(2,l)})x \in R_r(q')$ for any $x \in \mathcal{A}$, so we have

$$\begin{aligned} \text{dist}((1 - a\alpha_{p,q'}^{(2,l)})x, R_r(q)) &\leq \hat{\delta}(R_r(q'), R_r(q))\|(1 - a\alpha_{p,q'}^{(2,l)})x\| \\ &\leq \|q - q'\|\|(1 - a\alpha_{p,q'}^{(2,l)})x\|. \end{aligned}$$

Since $K_r(\alpha_{p,q}^{(2,l)}) = R_r(q)$, then for any $\epsilon > 0$, there is $z \in \mathcal{A}$ such that

$$\|(1 - \alpha_{p,q'}^{(2,l)})x - (1 - \alpha_{p,q}^{(2,l)})z\| \leq \|q - q'\| \|(1 - \alpha_{p,q'}^{(2,l)})x\| + \epsilon,$$

and then we have

$$\begin{aligned} \|(\alpha_{p,q'}^{(2,l)} - \alpha_{p,q}^{(2,l)})x\| &= \|\alpha_{p,q}^{(2,l)}(\alpha_{p,q'}^{(2,l)} - \alpha_{p,q}^{(2,l)})x\| \\ &\leq \|\alpha_{p,q}^{(2,l)}\| \|(1 - \alpha_{p,q'}^{(2,l)})x - (1 - \alpha_{p,q}^{(2,l)})z\| \\ &\leq (\|q - q'\| \|(1 - \alpha_{p,q'}^{(2,l)})x\| + \epsilon) \|\alpha_{p,q}^{(2,l)}\|. \end{aligned} \quad (4.8)$$

Since, we also have

$$\begin{aligned} \|(1 - \alpha_{p,q'}^{(2,l)})x\| &\leq \|x\| + \|a\| \|\alpha_{p,q}^{(2,l)}x - (\alpha_{p,q'}^{(2,l)} - \alpha_{p,q}^{(2,l)})x\| \\ &\leq (1 + \kappa)\|x\| + \|a\| \|(\alpha_{p,q'}^{(2,l)} - \alpha_{p,q}^{(2,l)})x\|. \end{aligned} \quad (4.9)$$

Now from Equations (4.8) and (4.9), we can compute

$$\|(\alpha_{p,q'}^{(2,l)} - \alpha_{p,q}^{(2,l)})x\| \leq \frac{(\|q - q'\| (1 + \kappa)\|x\| + \epsilon) \|\alpha_{p,q}^{(2,l)}\|}{1 - \kappa\|q - q'\|}.$$

Let $\epsilon \rightarrow 0^+$ in the above inequality, we obtain that

$$\frac{\|\alpha_{p,q'}^{(2,l)} - \alpha_{p,q}^{(2,l)}\|}{\|\alpha_{p,q}^{(2,l)}\|} \leq \frac{(1 + \kappa)\|q' - q\|}{1 - \kappa\|q' - q\|}, \quad \|\alpha_{p,q'}^{(2,l)}\| \leq \frac{1 + \|q' - q\|}{1 - \kappa\|q' - q\|} \|\alpha_{p,q}^{(2,l)}\|.$$

(2) By Lemma 4.5, $\alpha_{p,q'}^{(2,l)} = (va)^{(1,5)}v = v(av)^{(1,5)}$. For convenience, we write

$$b = \alpha_{p,q}^{(2,l)}, \quad b' = \alpha_{p,q'}^{(2,l)}, \quad \text{and} \quad x = \alpha_{p,q}^{(2,l)} + \alpha_{p,q}^{(2,l)}(av)^{(1,5)}a(v-w)(1 - \alpha_{p,q}^{(2,l)}).$$

Now we prove that $x = b'$ by using Lemma 2.2 (4). Obviously, we have

$$px = x. \quad \text{Note that } p = \alpha_{p,q}^{(2,l)}ap, \text{ so}$$

$$xap = \{\alpha_{p,q}^{(2,l)} + \alpha_{p,q}^{(2,l)}(av)^{(1,5)}a(v-w)(1 - \alpha_{p,q}^{(2,l)})\}ap$$

$$\begin{aligned}
 &= a_{p,q}^{(2,l)}ap + a_{p,q}^{(2,l)}(av)^{(1,5)}a(v-w)(ap-ap) \\
 &= p.
 \end{aligned}$$

Since $b = w(av)^{(1,5)}$ we have

$$\begin{aligned}
 xq' &= \{a_{p,q}^{(2,l)} + a_{p,q}^{(2,l)}(av)^{(1,5)}a(v-w)(1 - aa_{p,q}^{(2,l)})\}q' \\
 &= bq' + \{b(av)^{(1,5)}av - b(av)^{(1,5)}avab - b(av)^{(1,5)}aw + b(av)^{(1,5)}awab\}q' \\
 &= bq' + bab'q' - bab'abq' - b(av)^{(1,5)}awq' + b(av)^{(1,5)}awabq' \\
 &= bq' + 0 - bq' - b(av)^{(1,5)}awq' + b(av)^{(1,5)}awaw(av)^{(1,5)}q' \\
 &= 0.
 \end{aligned}$$

Thus, we have $x(1 - q') = x$. Finally, since $aw = awaw(av)^{(1,5)} = awab$, we have

$$\begin{aligned}
 (1 - q')ax &= (1 - q')a\{a_{p,q}^{(2,l)} + a_{p,q}^{(2,l)}(av)^{(1,5)}a(v-w)(1 - aa_{p,q}^{(2,l)})\} \\
 &= (1 - q')ab'a\{a_{p,q}^{(2,l)} + a_{p,q}^{(2,l)}(av)^{(1,5)}a(v-w)(1 - aa_{p,q}^{(2,l)})\} \\
 &= (1 - q')\{ab'ab + ab'ab(av)^{(1,5)}(av-w)(1 - ab)\} \\
 &= (1 - q')(ab + ab' - ab - ab'ab(av)^{(1,5)}aw + ab'ab(av)^{(1,5)}awab) \\
 &= (1 - q')ab' \\
 &= 1 - q'.
 \end{aligned}$$

Therefore, by Lemma 2.2 and the uniqueness of $a_{p,q}^{(2,l)}$, we get that

$$x = a_{p,q}^{(2,l)}. \quad \square$$

When the idempotents p and q both have some small perturbations, we have the following result:

Theorem 4.7. *Let $a \in \mathcal{A}$ and $p, q, p', q' \in \mathcal{A}^\bullet$ such that $a_{p,q}^{(2,l)}$ exists. If $\|p - p'\| < \frac{1}{(1+\kappa)^2}$ and $\|q - q'\| < \frac{1}{3+\kappa}$. Then $a_{p',q'}^{(2,l)}$ exists and*

$$\frac{\|a_{p',q'}^{(2,l)} - a_{p,q}^{(2,l)}\|}{\|a_{p,q}^{(2,l)}\|} \leq \frac{(1+\kappa)(\|p - p'\| + \|q - q'\|)}{1 - (1+\kappa)\|p - p'\| - \kappa\|q - q'\|};$$

$$\|a_{p',q'}^{(2,l)}\| \leq \frac{(1 + \|q - q'\|)\|a_{p,q}^{(2,l)}\|}{1 - (1+\kappa)\|p - p'\| - \kappa\|q - q'\|}.$$

Proof. By Theorem 4.4, $a_{p',q'}^{(2,l)}$ exists when $\|p - p'\| < \frac{1}{(1+\kappa)^2}$ and in this case

$$\|a_{p',q'}^{(2,l)}\| \leq \frac{\|a_{p,q}^{(2,l)}\|}{1 - (1+\kappa)\|p' - p\|} \leq \frac{1+\kappa}{\kappa} \|a_{p,q}^{(2,l)}\|.$$

So $\|q - q'\| < \frac{1}{3+\kappa} < \frac{1}{2 + \|a\|\|a_{p',q}^{(2,l)}\|}$ and consequently, $a_{p',q'}^{(2,l)}$ exists by

Theorem 4.6. Finally, by Theorems 4.4 and 4.6, we have

$$\begin{aligned} \|a_{p',q'}^{(2,l)} - a_{p,q}^{(2,l)}\| &\leq \|a_{p',q'}^{(2,l)} - a_{p',q}^{(2,l)}\| + \|a_{p',q}^{(2,l)} - a_{p,q}^{(2,l)}\| \\ &\leq \frac{(1 + \|a\|\|a_{p',q}^{(2,l)}\|)\|q - q'\|}{1 - \|a\|\|a_{p',q}^{(2,l)}\|\|q - q'\|} \|a_{p',q}^{(2,l)}\| + \frac{(1+\kappa)\|p - p'\|}{1 - (1+\kappa)\|p - p'\|} \|a_{p',q}^{(2,l)}\| \\ &\leq \frac{(1+\kappa)(1 - \|p - p'\|)\|q - q'\|}{1 - (1+\kappa)\|p - p'\| - \kappa\|q - q'\|} \frac{\|a_{p',q}^{(2,l)}\|}{1 - (1+\kappa)\|p - p'\|} \\ &\quad + \frac{(1+\kappa)\|p - p'\|}{1 - (1+\kappa)\|p - p'\|} \|a_{p',q}^{(2,l)}\| \\ &= \frac{(1+\kappa)(\|p - p'\| + \|q - q'\|)}{1 - (1+\kappa)\|p - p'\| - \kappa\|q - q'\|} \|a_{p,q}^{(2,l)}\|, \end{aligned}$$

$$\text{and } \|a_{p',q'}^{(2,l)}\| \leq \|a_{p',q'}^{(2,l)} - a_{p,q}^{(2,l)}\| + \|a_{p,q}^{(2,l)}\| \leq \frac{(1 + \|q - q'\|) \|a_{p,q}^{(2,l)}\|}{1 - (1 + \kappa) \|p - p'\| - \kappa \|q - q'\|}. \quad \square$$

Now, we consider the case when the elements $a, p, q \in \mathcal{A}^\bullet$ all have some small perturbations.

Theorem 4.8. *Let $a, \delta a \in \mathcal{A}$ and $p, q, p', q' \in \mathcal{A}^\bullet$ such that $a_{p,q}^{(2,l)}$ exists. If $\|p - p'\| < \frac{1}{(\kappa + 1)^2}$, $\|q - q'\| < \frac{1}{\kappa + 3}$, and $\|a_{p,q}^{(2,l)}\| \|\delta a\| < \frac{2\kappa}{(\kappa + 1)(\kappa + 4)}$.*

Then $\bar{a}_{p',q'}^{(2,l)}$ exists and

$$\begin{aligned} \|\bar{a}_{p',q'}^{(2,l)}\| &\leq \frac{(1 + \|q - q'\|) \|a_{p,q}^{(2,l)}\|}{1 - (1 + \kappa) \|p - p'\| - \kappa \|q - q'\| - (1 + \|q - q'\|) \|a_{p,q}^{(2,l)}\| \|\delta a\|}; \\ \|\bar{a}_{p',q'}^{(2,l)} - a_{p,q}^{(2,l)}\| &\leq \frac{\|a_{p,q}^{(2,l)}\|}{1 - (1 + \kappa) \|p - p'\| - \kappa \|q - q'\|} \\ &\left[(1 + \kappa) (\|p - p'\| + \|q - q'\|) + \frac{(1 + \|q - q'\|)^2 \|\delta a\| \|a_{p,q}^{(2,l)}\|}{1 - (1 + \kappa) \|p - p'\| - \kappa \|q - q'\| - (1 + \|q - q'\|) \|a_{p,q}^{(2,l)}\| \|\delta a\|} \right]. \end{aligned}$$

Proof. Theorem 4.7 indicates that $a_{p',q'}^{(2,l)}$ exists and

$$\|a_{p',q'}^{(2,l)}\| \leq \frac{(1 + \|q - q'\|) \|a_{p,q}^{(2,l)}\|}{1 - (1 + \kappa) \|p - p'\| - \kappa \|q - q'\|} < \frac{(1 + \kappa)(4 + \kappa)}{2\kappa} \|a_{p,q}^{(2,l)}\|.$$

Thus, $\|a_{p',q'}^{(2,l)}\| \|\delta a\| < 1$ and hence $1 + a_{p',q'}^{(2,l)} \delta a$ is invertible. Therefore, $a_{p',q'}^{(2,l)}$ exists and $\bar{a}_{p',q'}^{(2,l)} = a_{p',q'}^{(2,l)} (1 + \delta a a_{p',q'}^{(2,l)})^{-1}$ by Theorem 3.4. Now by Theorem 4.7, we have

$$\begin{aligned} \|\bar{a}_{p',q'}^{(2,l)}\| &\leq \frac{\|a_{p',q'}^{(2,l)}\|}{1 - \|a_{p',q'}^{(2,l)}\| \|\delta a\|} \\ &\leq \frac{[1 + \|q - q'\|] \|a_{p,q}^{(2,l)}\|}{1 - [1 + \kappa] \|p - p'\| - \kappa \|q - q'\| - [1 + \|q - q'\|] \|a_{p,q}^{(2,l)}\| \|\delta a\|}, \end{aligned}$$

and

$$\begin{aligned}
\|\bar{\alpha}_{p',q'}^{(2,l)} - \alpha_{p,q}^{(2,l)}\| &\leq \|(1 + \alpha_{p',q'}^{(2,l)}\delta\alpha)^{-1}\alpha_{p',q'}^{(2,l)} - \alpha_{p',q'}^{(2,l)}\| + \|\alpha_{p',q'}^{(2,l)} - \alpha_{p,q}^{(2,l)}\| \\
&\leq \frac{\|\delta\alpha\|\|\alpha_{p',q'}^{(2,l)}\|^2}{1 - \|\delta\alpha\|\|\alpha_{p',q'}^{(2,l)}\|} + \|\alpha_{p',q'}^{(2,l)} - \alpha_{p,q}^{(2,l)}\| \\
&\leq \frac{\|\alpha_{p,q}^{(2,l)}\|}{1 - (1 + \kappa)\|p - p'\| - \kappa\|q - q'\|} \\
&\left[(1 + \kappa)(\|p - p'\| + \|q - q'\|) + \frac{(1 + \|q - q'\|)^2\|\delta\alpha\|\|\alpha_{p,q}^{(2,l)}\|}{1 - (1 + \kappa)\|p - p'\| - \kappa\|q - q'\| - (1 + \|q - q'\|)\|\alpha_{p,q}^{(2,l)}\|\|\delta\alpha\|} \right].
\end{aligned}$$

This completes the proof. \square

By using perturbation theorems for the generalized inverse $\alpha_{p,q}^{(2,l)}$, we can also investigate the perturbation analysis for the generalized inverse $\alpha_{p,q}^{(1,2)}$ under some conditions.

Corollary 4.9. *Let $a \in \mathcal{A}$ and $p, q \in \mathcal{A}^\bullet$ such that $\alpha_{p,q}^{(1,2)}$ exists.*

Suppose that $p' \in \mathcal{A}^\bullet$ with $\|p - p'\| < \frac{1}{(1 + \kappa)^2}$ and $ap' = a$. Then $\alpha_{p',q}^{(1,2)}$

exists and

$$\frac{\|\alpha_{p',q}^{(1,2)} - \alpha_{p,q}^{(1,2)}\|}{\|\alpha_{p,q}^{(1,2)}\|} \leq \frac{(1 + \kappa)\|p' - p\|}{1 - (1 + \kappa)\|p' - p\|} \text{ and } \|\alpha_{p',q}^{(1,2)}\| \leq \frac{\|\alpha_{p,q}^{(1,2)}\|}{1 - (1 + \kappa)\|p' - p\|}.$$

Proof. Set $b = \alpha_{p,q}^{(1,2)}$. Then $bab = b$, $aba = a$, $ba = p$, $1 - ab = q$,

and $\alpha_{p',q}^{(2,l)} = \alpha_{p,q}^{(1,2)} = b$. By Theorem 4.4, $\alpha_{p',q}^{(2,l)}$ is exists and

$$\frac{\|\alpha_{p',q}^{(2,l)} - \alpha_{p,q}^{(1,2)}\|}{\|\alpha_{p,q}^{(1,2)}\|} \leq \frac{(1 + \kappa)\|p' - p\|}{1 - (1 + \kappa)\|p' - p\|} \text{ and } \|\alpha_{p',q}^{(2,l)}\| \leq \frac{\|\alpha_{p,q}^{(1,2)}\|}{1 - (1 + \kappa)\|p' - p\|}.$$

We need only to show that $\alpha_{p',q}^{(1,2)} = \alpha_{p',q}^{(2,l)}$ in this case. Put $b' = \alpha_{p',q}^{(2,l)}$. Then $b'ab' = b'$, $(b'a)\mathcal{A} = p'\mathcal{A}$, $(1 - ab')\mathcal{A} = q\mathcal{A}$. Thus $(1 - q)(1 - ab') = 0$ and hence $1 - q = (1 - q)ab' = abab' = ab'$. Furthermore, $ab'a = (1 - q)a = aba = a$. From $(b'a)\mathcal{A} = p'\mathcal{A}$, we get that $(1 - b'a)p' = 0$ and $p' = b'ap' = b'a$. Therefore, $b' = \alpha_{p',q}^{(1,2)}$. \square

We need the following easy representation lemma for $\alpha_{p,q}^{(1,2)}$:

Lemma 4.10. *Let $a \in \mathcal{A}$ and $p, q \in \mathcal{A}^\bullet$ such that $\alpha_{p,q}^{(1,2)}$ exists. Let $w \in \mathcal{A}$ such that $wa = p$ and $aw = 1 - q$. Then $\alpha_{p,q}^{(1,2)} = (wa)^\# w = w(aw)^\#$.*

Proof. Obviously, $wa, aw \in \mathcal{A}^g$ for $wa = p$ and $aw = 1 - q$. We also have $(wa)^\# = p$ and $(aw)^\# = 1 - q$. Then by using the uniqueness of $\alpha_{p,q}^{(1,2)}$, we can prove our lemma by simple computation. \square

Corollary 4.11. *Let $a \in \mathcal{A}$ and $p, q \in \mathcal{A}^\bullet$ such that $\alpha_{p,q}^{(1,2)}$ exists. Suppose that $q' \in \mathcal{A}^\bullet$ with $\|q - q'\| < \frac{1}{2 + \kappa}$ and $a = (1 - q')a$. Then $\alpha_{p,q'}^{(1,2)}$ exists and*

$$(1) \frac{\|\alpha_{p,q'}^{(1,2)} - \alpha_{p,q}^{(1,2)}\|}{\|\alpha_{p,q}^{(1,2)}\|} \leq \frac{(1 + \kappa)\|q - q'\|}{1 - \kappa\|q - q'\|} \text{ and } \|\alpha_{p,q'}^{(1,2)}\| \leq \frac{1 + \|q' - q\|}{1 - \kappa\|q' - q\|} \|\alpha_{p,q}^{(1,2)}\|.$$

(2) *If there are some $w, v \in \mathcal{A}$ with $wa = p = va$, $aw = 1 - q$, and $av = 1 - q'$. Then*

$$\alpha_{p,q'}^{(1,2)} = \alpha_{p,q}^{(1,2)} + \alpha_{p,q}^{(1,2)}(av)^\# a(v - w)q.$$

Proof. $\alpha_{p,q'}^{(2,l)}$ exists by Theorem 4.6. From $a = (1 - q')a$ and Lemma 2.2, we can obtain that $\alpha_{p,q'}^{(1,2)}$ exists and $\alpha_{p,q'}^{(2,l)} = \alpha_{p,q'}^{(1,2)}$.

Now, the estimates in (1) and the representation for $\alpha_{p,q}^{(1,2)}$ in (2) follow from Theorem 4.6 and Lemma 4.10. \square

Finally, by Corollaries 4.9, 4.11, and Theorem 4.7, we have

Corollary 4.12. *Let $a \in \mathcal{A}$ and $p, q, p', q' \in \mathcal{A}^\bullet$ with $\alpha_{p,q}^{(1,2)}$ exists.*

If $\|p - p'\| < \frac{1}{(1 + \kappa)^2}$, $\|q - q'\| < \frac{1}{3 + \kappa}$, and $ap' = a = (1 - q')a$. Then

$\alpha_{p',q'}^{(1,2)}$ exists and

$$\frac{\|\alpha_{p',q'}^{(1,2)} - \alpha_{p,q}^{(1,2)}\|}{\|\alpha_{p,q}^{(1,2)}\|} \leq \frac{(1 + \kappa)(\|p - p'\| + \|q - q'\|)}{1 - (1 + \kappa)\|p - p'\| - \kappa\|q - q'\|};$$

$$\|\alpha_{p',q'}^{(1,2)}\| \leq \frac{(1 + \|q - q'\|)\|\alpha_{p,q}^{(1,2)}\|}{1 - (1 + \kappa)\|p - p'\| - \kappa\|q - q'\|}.$$

References

- [1] J. Cao and Y. Xue, The characterizations and representations for the generalized inverses with prescribed idempotents in Banach algebras, (submitted).
- [2] G. Chen, Y. Wei and Y. Xue, Perturbation analysis of the least square solution in Hilbert spaces, *Linear Algebra Appl.* 244 (1996), 69-80.
- [3] G. Chen and Y. Xue, Perturbation analysis for the operator equation $Tx = b$ in Banach spaces, *J. Math. Anal. Appl.* 212 (1997), 107-125.
- [4] G. Chen and Y. Xue, The expression of the generalized inverse of the perturbed operator under type I perturbation in Hilbert spaces, *Linear Algebra Appl.* 285 (1998), 1-6.
- [5] D. Cvetkovic-Ilić, X. Liu and J. Zhong, On the (p, q) -outer generalized inverse in Banach Algebras, *Appl. Math. Comput.* 209(2) (2009), 191-196.
- [6] D. S. Djordjević and Y. Wei, Outer generalized inverses in rings, *Comm. Algebra* 33 (2005), 3051-3060.
- [7] F. Du and Y. Xue, Perturbation analysis of $A_{T,S}^{(2)}$ on Banach spaces, *Electronic J. Linear Algebra* 23 (2012), 586-598.

- [8] F. Du and Y. Xue, The perturbation of the group inverse under the stable perturbation in a unital ring, *Filomat* 27 (2013), 61-70.
- [9] T. Kato, *Perturbation Theory for Linear Operators*, Springer-Verlag, New York, 1984.
- [10] V. Müller, *Spectral Theory of Linear Operators and Spectral Systems in Banach Algebras*, 2nd Edition, Birkhäuser Verlag AG, 2007.
- [11] M. Z. Nashed, Inner, outer and generalized inverses in Banach and Hilbert spaces, *Numer. Funct. Anal. Optim.* 9 (1987), 261-325.
- [12] Y. Xue, Stable perturbation in Banach spaces, *J. Aust. Math. Soc.* 83 (2007), 1-14.
- [13] Y. Xue, *Stable Perturbations of Operators and Related Topics*, World Scientific, 2012.
- [14] Y. Xue and G. Chen, Some equivalent conditions of stable perturbation of operators in Hilbert spaces, *Appl. Math. Comput.* 147 (2004), 765-772.
- [15] Q. Xu, Y. Wei and Y. Gu, Sharp norm-estimation for Moore-Penrose inverses of stable perturbations of Hilbert C^* -module operators, *SIAM J. Numer. Anal.* 47 (2010), 4735-4758.

